

European Journal of Radiology

Volume 81, Issue 4, April 2012, Pages e561–e564

Magnetic resonance imaging of the lumbar spine with axial loading: A review of 120 cases

Andre Kinder^{a,b,1}, Fernando Palma Filho^{b,2}, Elisio Ribeiro^{b,2}, Romeu C. Domingues^{b,4}, Roberto C. Domingues^{b,4}, Edson Marchiori^{a,5}, Emerson Gasparetto^{a,b}.

^a Department of Radiology, Federal University of Rio de Janeiro School of Medicine, Rio de Janeiro, Brazil

^b Clinica de Diagnóstico por Imagem (CDPI), Rio de Janeiro, Brazil

Received 14 March 2011. Accepted 7 June 2011. Available online 8 July 2011.

Abstract

Purpose

To evaluate the imaging findings of patients with clinical symptoms of lower back pain who underwent magnetic resonance imaging (MRI) of the lumbar spine with axial loading.

Materials and methods

We examined 120 patients by MRI, before and after axial loading, using a compression device that applied 50% of their body weight for a load time of 5 min. The dural sac cross area (DSCA) was examined by two experienced radiologists before and after axial load, and their findings were compared. Degenerative abnormalities within and adjacent to the spinal canal were also analyzed.

Results

A reduction in DSCA greater than 15 mm² after axial load was defined as significant, and was found in 81 patients (67.5%) and 138 disc spaces (38.3%). Reduction was most frequent at L4-L5 ($n = 55$). For other disorders, a 9% increase in cases of bulging disc was seen during axial loading, and seven disc spaces showed protrusion/extrusion only after load. Facet joint synovial cysts, foraminal stenosis, and hypertrophy of the flavum ligaments showed almost no differences, pre- and post-load.

Conclusion

For adequate evaluation of lumbar symptoms, examination should be performed with axial loading, especially in cases of suspected spinal stenosis.

Keywords

Magnetic resonance imaging; Lumbar spine; Axial loading; Spinal stenosis

Figures and tables from this article:




Fig. 1. Reduction of DSCA at normal L4-L5 disc level. (A) Axial T2-weighted spin-echo sequence without (A) and with (B) axial loading. DSCA was 115 mm² before loading and 88 mm² after that. In addition to a posterior disc protrusion, the post-loading image demonstrates hypertrophy of the ligamentum flavum.




Fig. 2. Development of occult facet joint cyst. (A) Axial T2-weighted fast spin-echo sequence without axial loading. (B) Axial T2-weighted spin-echo sequence with axial loading shows a right facet joint cyst deep in the ligamentum flavum, contributing to the central stenosis.

Corresponding author at: Hospital Universitário Clementino Fraga Filho/UFRJ, Rua Professor Rodolpho Paulo Rocco, no. 255, Departamento de Radiologia, Subsolo, CEP: 21941-617 Ilha do Fundão, Rio de Janeiro, RJ, Brazil. Tel.: +55 21 2562 2568; fax: +55 21 2562 2568.

1. Contact address: Rua Dr. Nelson de Sá Earp, 144, 401, Centro, Petrópolis, CEP: 25680-195, Brazil.
2. Contact address: Rua Saddock de Sá, 266, Ipanema, CEP: 22411-040 Rio de Janeiro, Brazil.
3. Contact address: Av. Rainha Elizabeth 371 ap 607, Copacabana, CEP: 22081-031 Rio de Janeiro, Brazil.
4. Contact address: Av. das Américas, 4666, grupos 302A, Barra da Tijuca, CEP: 22790-972 Rio de Janeiro, Brazil.
5. Contact address: Rua Thomaz Cameron, 438, Valparaíso, CEP: 25685-120 Petrópolis, Rio de Janeiro, Brazil.